


CLAIMS:

1. A compound having the structure of Formula I



FORMULA I

and its pharmaceutically acceptable salts, enantiomers, diastereomers, N-oxides, prodrugs or metabolites, wherein

T is five to seven membered heterocyclic ring, aryl, substituted aryl, bound to the ring C with a linker W and the heterocyclic and aryl rings are further substituted by a group represented by R,

wherein R is selected from the group consisting of alkyl (C₁₋₆), halogen-CN, COR₅, COOR₅, N(R₆, R₇), CON (R₆, R₇), CH₂NO₂, NO₂, CH₂R₈, CHR₉, -CH = N-OR₁₀, -C=CH-R₅, wherein R₅ is selected from the group consisting of H, optionally substituted C₁-C₁₂, alkyl, C₃₋₁₂, cycloalkyl, aryl, heteroaryl; R₆ and R₇ are independently selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₆ alkoxy; R₈ and R₉ are independently selected from the group consisting of H, C₁₋₆ alkyl, F, Cl, Br, C₁₋₁₂ alkyl substituted with one or more of F, Cl, Br, I, OR₄, SR₄, N(R₆, R₇) wherein R₄ is selected from the group consisting of H, C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl substituted with one or more F, Cl, Br, I or

OH and R_6 and R_7 are the same as defined earlier, R_{10} is selected from the group consisting of H, optionally substituted from H, optionally substituted C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl, aryl, heteroaryl;

n is an integer in the range from 0 to 3;

5 **X** is CH, CH-S, CH-O and N


Y and Z are independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{3-12} cycloalkyl, C_{0-3} bridging group;

U and V are independently selected from the group consisting of optionally substituted C_{1-6} alkyl, F, Cl, Br, C_{1-12} alkyl substituted with one or more of F, Cl, Br, I, preferably U and V are hydrogen or fluoro;

W is selected from the group CH_2 , CO, CH_2NH , $-NHCH_2$, $-CH_2NHCH_2$, $-CH_2-N(R_{11})CH_2-$, $-CO-CO-$, $CH_2(R_{11})N-$, $CH(R_{11})$, S, $CH_2(CO)$, N(R_{11}) wherein R_{11} is hydrogen, optionally substituted C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl, aryl or heteroaryl;

15 **R₁** is selected from the group consisting of $-NHC(=O)R_2$ wherein R_2 is hydrogen, C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl substituted with one or more of F, Cl, Br, I or OH; $N(R_3, R_4)$; $-NR_2C(=S)R_3$; $-NR_2C(=S)SR_3$ wherein R_2 is the same as defined above and R_3 and R_4 are independently selected from the group consisting of H, C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl substituted with one or more of F, Cl, Br, I or OH.

2. A compound having structure of Formula II



FORMULA II

and its pharmaceutically acceptable salts, enantiomers, diastereomers, N-oxides, prodrugs or metabolites wherein

M= O, S, NH, N-CH₃;

X is CH, CH-S, CH-O and N;


Y and Z are independently selected from the group consisting of hydrogen , C₁₋₆ alkyl, C₃₋₁₂ cycloalkyl, C₀₋₃ bridging group;

U and V are independently selected from the group consisting of optionally substituted C₁₋₆ alkyl, F, Cl, Br, C₁₋₁₂ alkyl substituted with one or more of F, Cl, Br, I, preferably U and V are hydrogen or fluoro;


W is selected from the group consisting of CH₂, CO, CH₂NH, -NHCH₂, -CH₂NHCH₂, -CH₂-N (R₁₁) CH₂ -, CH₂ (R₁₁) N -, CH (R₁₁) , S, CH₂(CO), NH wherein R₁₁ is optionally substituted with C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl , aryl , heteroaryl except when M=S, Q=P=H, W=(C=O);

n is an integer in the range from 0 to 3; and,


Q and P are independently selected from the group consisting of $-\text{CN}$, COR_5 , COOR_5 , $\text{N}(\text{R}_6, \text{R}_7)$, $\text{CON}(\text{R}_6, \text{R}_7)$, CH_2NO_2 , NO_2 , CH_2R_8 , CHR_9 , $-\text{CH}=\text{N}-\text{OR}_{10}$, $\text{C}=\text{CH}-\text{R}_5$, wherein R_5 is selected from the group consisting of H , optionally substituted C_{1-12} alkyl, C_{3-12} cycloalkyl, aryl, heteroaryl; R_6 and R_7 are independently selected from the group consisting of H , optionally substituted C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy; R_8 and R_9 are independently selected from the group consisting of H , C_{1-6} alkyl, F , Cl , Br , C_{1-12} alkyl substituted with one or more of F , Cl , Br , I , OR_4 , SR_4 , wherein R_4 is selected from the group consisting of H , C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl substituted with one or more F , Cl , Br , I or OH , $\text{N}(\text{R}_6, \text{R}_7)$, R_{10} is selected from the group consisting of H , optionally substituted C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl, aryl, heteroaryl except $\text{W}=(\text{CO})$, Q and $\text{P}=\text{H}$ and $\text{M}=\text{S}$, ring C in Formula II is 6-8 membered or of larger size and the larger rings have either two or three carbons between each nitrogen atom, comprising of



and may be bridged to form a bicyclic system as shown below,




ring C is optionally substituted by Y and Z with alkyl groups, cycloalkyl groups, fluoro group, carboxylic and corresponding esters, amides, substituted alkyls or bridging alkyl groups are as shown below:



six membered ring C with $X = -CH-(NR_{11})$, (wherein R_{11} is the same as defined earlier) is selected from the group consisting of the following rings;

5




10



15




20




25

30




35

40




45

wherein M = Sulphur and Oxygen as shown by Formulae III and IV respectively,



Formula III



Formula IV

wherein P, Q, U, V, X, Y, Z, W and n in Formulae III and IV as defined earlier for Formula I.

3. A compound selected from the group consisting of
 1. (S)-N-[[3-[3-Fluoro-4-[N-1-[4-(2-furoyl) piperazinyl]]phenyl]-2-oxo-5-oxazolidinyl] methyl]acetamide
 2. (S)-N-[[3-[3-Fluoro-4-[N-1[4-{2-furyl(5-formyl)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide

3. (S)-N-[[3-Fluoro-4-[N-1[4-(2-furyl)-(5-carboxyethyl)methyl]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
- 5 4. (S)-N-[[3-Fluoro-4-[N-1[4-(5-bromo-2-furoyl)]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl] methyl]acetamide
5. (S)-N-[[3-Fluoro-4-[N-1[4-(5-chloromethyl-2-furoyl)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
6. (S)-N-[[3-Fluoro-4-[N-1[4-(5-nitro-2-furoyl)piperazinyl]phenyl]-2-oxo-5-oxazolidinyl] methyl]acetamide
- 10 7. (S)-N[[3-[3-Fluoro-4-[N-1[4-{2-(2-thienyl)dicarbonyl}]piperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl]acetamide
8. (S)-N[[3-[3-Fluoro-4-[N-1[4-(3-furoyl)]piperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl] acetamide
- 15 9. (S)-N[[3-[3-Fluoro-4-[N-1[4-{2-furyl(5-bromo)methyl}]piperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl]acetamide
10. (S)-N[[3-[3-Fluoro-4-[N-1[4-{2-thienyl(5-chloro)methyl}]piperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl]acetamide
- 20 11. (S)-N[[3-[3-Fluoro-4-[N-1[4-(2-furylmethyl)]piperazinyl]phenyl]2-oxo-5-oxazolidinyl] methyl]acetamide
12. (S)-N-[[3-[3-Fluoro-4-[N-1[4-(2-thienylmethyl)]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
- 25 13. (S)-N[[3-[3-Fluoro-4-[N-1[4-(2-thienylacetyl)]piperazinyl]phenyl]2-oxo-5-oxazolidinyl] methyl]acetamide
14. (S)-N-[[3-[3-Fluoro-4-[N-1[4-{2-thienyl(4-bromo)methyl}]piperazinyl]phenyl]-2 oxo-5-oxazolidinyl]methyl]acetamide
- 30 15. (S)-N-[[3-[3-fluoro-4-[N-1-[4-{2-furyl-(5-nitro)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide.

16. Hydrochloric salt of (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-nitro)methyl}]piperazinyl] phenyl]-2-oxo-5-oxazolidinyl)methyl]acetamide
- 5 17. Citrate salt of (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-nitro)methyl}]piperazinyl] phenyl]-2-oxo-5-oxazolidinyl)methyl]acetamide
18. (S)-N[[3-[3-Fluoro-4-[N-1[4-(2-pyrrolylmethyl)]piperazinyl]phenyl]2-oxo-5-oxazolidinyl)methyl]acetamide
- 10 19. (S)-N[[3-[3-Fluoro-4-[N-1[4-{2-thienyl(3-methyl)methyl}]piperazinyl]phenyl]2-oxo-5-oxazolidinyl)methyl]acetamide
20. (S)-N[[3-[3-Fluoro-4-[N-1[4-(3-furylmethyl)]piperazinyl]phenyl]2-oxo-5-oxazolidinyl] methyl]acetamide
- 15 21. (S)-N[[3-[3-Fluoro-4-[N-1[4-{2-thienyl(5-methyl)methyl}]piperazinyl]phenyl]2-oxo-5-oxazolidinyl)methyl]acetamide
22. (S)-N[[3-[3-Fluoro-4-[N-1[4-{2-pyrrole(1-methyl)methyl}]piperazinyl] phenyl]2-oxo-5-oxazolidinyl)methyl]acetamide
- 20 23. (S)-N[[3-[3-Fluoro-4-[N-1[4-{2-thienyl(5-nitro)methyl}]piperazinyl]phenyl]2-oxo-5-oxazolidinyl)methyl]acetamide
- 25 24. (S)-N[[3-[3-Fluoro-4-[N-1[4-[2-furyl{5-(N-thiomorpholinyl)methyl}methyl]piperazinyl] phenyl]2-oxo-5-oxazolidinyl)methyl]acetamide
- 25 25. (S)-N[[3-[3-Fluoro-4-[N-1[4-[2-furyl{5-(N-morpholinyl)methyl}methyl]]piperazinyl] phenyl]2-oxo-5-oxazolidinyl)methyl]acetamide
26. (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-acetoxymethyl)methyl}]piperazinyl] phenyl]-2-oxo-5-oxazolidinyl)methyl]acetamide
- 30 27. (S)-N-[[3-Fluoro-4-[N-1[4-{2-thienyl(5-bromo)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl)methyl]acetamide
28. (S)-N-[[3-Fluoro-4-[N-1[4-(5-nitro-2-furylmethyl)piperazinyl] phenyl]- 2-oxo oxazolidinyl)methyl]dichloroacetamide

29. (S)-N-[[3-[3-Fluoro-4-[N-1[4-(5-nitro-2-thienoyl)]piperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl]acetamide hydrochloride
30. (S)-N-[[3-[3-Fluoro-4-[N-1[4-(2',2'- diphenyl-2' hydroxy acetyl)]piperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl]acetamide
- 5 31. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(5-nitro-2-furoyl)-N-methyl]amino]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
32. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(3-furoyl)-N-methyl]amino]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
- 10 33. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(5-bromo -2-furoyl)-N-methyl]amino]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
34. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(5-nitro-2-thienylmethyl)-N-methyl]amino]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
- 15 35. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(5-nitro-2-furylmethyl)-N-methyl] amino]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
36. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(5-formyl-2-furylmethyl)-N-methyl] amino-methyl]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl] methyl]acetamide
- 20 37. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(5-carboxyethyl-2-furylmethyl)-N-methyl] aminomethyl]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl] acetamide
- 25 38. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(2-thiopheneacetyl)-N-methyl]aminomethyl]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
39. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(5-nitro-2-thienylmethyl)-N-methyl]amino-methyl]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
- 30

40. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(5-nitro-2-furylmethyl)-N-methyl]amino-methyl]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
- 5 41. (S)-N-[[3-[4-[4-(N-methyl-N-2furyl(5formyl)methylaminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl]methyl]acetamide
42. (S)-N-[[3-[4-[4-(N-methyl-N-(3,5-difluorobenzoyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl] methyl]acetamide.
43. (S)-N-[[3-[4-[4-(N-methyl-N-(5-bromo-2-furoyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl] methyl]acetamide
- 10 44. (S)-N-[[3-[4-[4-(N-methyl-N-(5-nitro-2-furoyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl]methyl]acetamide
45. (S)-N-[[3-[4-[4-(N-methyl-N-3- furoyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl]methyl]acetamide.
- 15 46. (S)-N-{3-[4-[4-(N-methyl, N- 2-furoyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl methyl]acetamide
47. (S)-N-{3-[4-[4-(N-methyl,2-thiopheneacetyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo oxazolidin-5-yl methyl]acetamide
48. (S)-N-[[3-[4-[4-(N-methyl-N-2furylmethyl) aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl]methyl]acetamide
- 20 49. (S)-N-[[3-[4-[4-(N-methyl-N-3-furyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl] methyl]acetamide.
50. (S)-N-[[3-[4-[4-(N-methyl-N-2-furyl(5-nitro)methyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl] methyl]acetamide.
- 25 51. (S)-N-[[3-[4-[4-(N-methyl-N-2-thienyl(5-nitro)methyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl] methyl]acetamide.
52. (S)-N-[[3-[4-[4-(N-methyl-N-2-thienylmethyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl]methyl]acetamide.
53. (S)-N-[[3-[4-[4-(N-methyl-N-(5-methyl-2-thienylmethyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl] methyl]acetamide
- 30 54. (S)-N-{3-[4-[4-(N-methyl,2-(5-bromo)thienylmethyl)aminopiperidine-1-yl]-3-fluorophenyl]-2-oxo-oxazolidin-5-yl methyl]acetamide

55. (S)-N-[[3-[3-Fluoro-4-[N-1[4-{2-furyl(5-formyl)methyl}]homopiperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl]acetamide
56. (S)-N-[[3-[3-Fluoro-4-[N-1[4-(2-thienylacetyl)]homopiperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl]acetamide
57. (S)-N-[[3-[3-Fluoro-4-[N-1[4-{2-thienyl(5-nitro)methyl}]homopiperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl]acetamide
58. (S)-N-[[3-[3-Fluoro-4-[N-1[4-(3-furylmethyl)]homopiperazinyl]phenyl]2-oxo-5-oxazolidinyl]methyl]acetamide
59. Preparation of (S)-N-[[3-[3-fluoro-4-[N-1{2-furyl-[4-(5-difluoromethyl)methyl]}piperazinyl]-2-oxo-5-oxazolidinyl]-methyl]acetamide.
60. (S)-N-[[3-[3-Fluoro-4-[N-1-[4-(2-furyl(5-aldoxime)methyl)]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
61. (S)-N-[[3-[3-Fluoro-4-[N-1[4-{2-furyl(5-aldoxime(methyl-4-(N-carboxyamino)phenyl acetate) methyl]}piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
62. (S)-N-[[3-[3-Fluoro-4[N-1-[4-{2-furyl(5-hydrazone)-methyl}]piperazinyl]-phenyl]-2-oxo-5-oxazolidinyl]-methyl]acetamide
63. Preparation of (S)-N-[[3-[3-Fluoro-4-[N-1{2-furyl-[4-(5-hydroxymethyl)methyl]}piperazinyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
64. (S)-N-[[3-[3-Fluoro-4-[N-1[4-{2-furyl(5-cyano)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
65. (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-carboxy)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
66. (S)-N-[[3-Fluoro-4-[N-1[5-(1,3-dioxane)-2-furylmethyl]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
67. (S)-N-[[3-Fluoro-4-[N-1[5-(formamido)-2-furylmethyl]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide
68. (S)-N-[[3-Fluoro-4-[N-1[5-(morpholine-1-carbonyl)-2-furylmethyl]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide

69. (S)-N-[[3-Fluoro-4-[N-1[5-(4-(tert butoxy carbonyl)amino piperidine)-2-furylmethyl]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl] acetamide

70. (S)-N-[[3-Fluoro-4-[N-1[4-{(Z)-2-methoxyimino-2-(2-furyl)acetyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide

5 71. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(2-thiopheneacetyl)-N-methyl]amino]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide

72. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(5-formyl-2-furylmethyl)-N-methyl]amino]-3-azabicyclo-[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide

10 73. (S)-N-[[3-[3-Fluoro[4-[3-(1 α ,5 α ,6 α)-6-[N-(3-thienoyl)-N-methyl]amino]-3-azabicyclo[3.1.0]hexane]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide

15 74. (S)-N-[[3-[3-fluoro-4-[N-1{2-furyl-[4-(5-fluoromethyl)methyl}]piperazinyl]-2-oxo-5-oxazolidinyl]-methyl]acetamide.


4. A pharmaceutical composition comprising the compound of claims 1, 2, or 3 and a pharmaceutical acceptable carrier.

5. A pharmaceutical composition comprising a pharmaceutically effective amount of compound according to claims 1, 2, or 3, or a physiologically acceptable acid addition salt thereof with a pharmaceutical acceptable carrier for treating microbial infections.

20 6. A method of treating or preventing microbial infections in a mammal comprising administering to the said mammal, the pharmaceutical composition according to claim 5.

25

7. A process for preparing a compound of Formula I



FORMULA I

and its pharmaceutically acceptable salts, enantiomers, diastereomers, N-oxides, prodrugs or metabolites, wherein

T is five to seven membered heterocyclic ring, aryl, substituted aryl, bound to the ring C with a linker w and the heterocyclic and aryl rings are further substituted by a group represented by R,

wherein R is selected from the group consisting of $-\text{CN}$, COR_5 , COOR_5 , $\text{N}(\text{R}_6, \text{R}_7)$, $\text{CON}(\text{R}_6, \text{R}_7)$, CH_2NO_2 , NO_2 , CH_2R_8 , CHR_9 , $-\text{CH} = \text{N}-\text{OR}_{10}$, $-\text{C}=\text{CH}-\text{R}_5$, wherein R_5 is selected from the group consisting of H, optionally substituted C_{1-12} , alkyl, C_{3-12} , cycloalkyl, aryl, heteroaryl, R_6 and R_7 , are independently selected from the group consisting of H, optionally substituted C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy; R_8 and R_9 are independently selected from the group consisting of H, C_{1-6} alkyl, F, Cl, Br, C_{1-12} alkyl substituted with one or more of F, Cl, Br, I, OR_4 , SR_4 , $\text{N}(\text{R}_6, \text{R}_7)$ wherein R_4 is selected from the group consisting of H, C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl substituted with one or more F, Cl, Br, I or OH and R_6 and R_7 are the same as defined earlier, R_{10} is selected from the group consisting of

H, optionally substituted from H, optionally substituted C₁₋₁₂ alkyl, C₃₋₅₁₂ cycloalkyl, C₁₋₆, alkoxy, C₁₋₆ alkyl, aryl, heteroaryl;

n is an integer in the range from 0 to 3;

X is CH, CH-S, CH-O and N;


5 Y and Z are independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₃₋₁₂ cycloalkyl, C₀₋₃ bridging group;

U and V are independently selected from the group consisting of optionally substituted C₁₋₆ alkyl, F, Cl, Br, C₁₋₁₂ alkyl substituted with one or more of F, Cl, Br, I, preferably U and V are hydrogen or fluoro;

10 W is selected from the group consisting of CH₂, CO, CH₂NH, -NHCH₂, -CH₂NHCH₂, -CH₂-N(R₁₁)CH₂-, CH₂(R₁₁)N-, CH(R₁₁), S, CH₂(CO), NH wherein R₁₁ is optionally substituted with C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl, aryl, heteroaryl; and

15 R₁ is selected from the group consisting of -NHC(=O)R₂ wherein R₂ is hydrogen, C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl substituted with one or more of F, Cl, Br, I or OH; N(R₃, R₄); -NR₂C(=S)R₃; -NR₂C(=S)SR₃ wherein R₂ is the same as defined above and R₃ and R₄ are independently selected from the group consisting of H, C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl substituted with one or more of F, Cl, Br, I or OH,

which comprises reacting an amine compound of Formula V




FORMULA V

with a heterocyclic compound of Formula R-T-W- R₁₂ wherein G in amines of Formula V is defined as NH, CH(NHR₁₃), -CH-CH₂NHR₁₃ wherein R₁₃ is H, ethyl, methyl, isopropyl, acetyl, cyclopropyl, alkoxy or acetyl and Y, Z, U, V, R₁, n, R, T and W are the same as defined earlier and R₁₂ is a suitable leaving group selected from the group comprising of fluoro, chloro, bromo, SCH₃, -SO₂CH₃, -SO₂CF₃ or OC₆H₅.

8. A process for preparing a compound of Formula I as claimed in claim 7, wherein W=CH₂ and R-T-W-R₁₂ is a five membered heterocyclic ring with aldehyde group and the compound of Formula I is produced by reductive amination.
9. A process for preparing a compound of Formula I as claimed in claim 7, wherein W = CO and R-T-W-R₁₂ is a five membered heterocyclic ring with carboxylic acid, and amino compound of Formula V is acylated with activated esters in presence of condensing agents comprising 1,3-dicyclohexylcarbodiimide (DCC) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC).

10. A process for the preparation of compound of Formula II



FORMULA II

wherein

n is an integer in the range from 0 to 3;

X is CH, CH-S, CH-O and N;

Y and Z are independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{3-12} cycloalkyl, C_{0-3} bridging group;


U and V are independently selected from the group consisting of optionally substituted C_{1-6} alkyl, F, Cl, Br, C_{1-12} alkyl substituted with one or more of F, Cl, Br, I, preferably **U** and **V** are hydrogen or fluoro;

W is selected from the group consisting of CH_2 , CO, CH_2NH , $-NHCH_2$, $-CH_2NHCH_2$, $-CH_2-N(R_{11})CH_2-$, $CH_2(R_{11})N-$, $CH(R_{11})$, S, $CH_2(CO)$, NH wherein R_{11} is optionally substituted with C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl, aryl, heteroaryl; and


Q and P are independently selected from the group consisting of $-CN$, COR_5 , $COOR_5$, $N(R_6, R_7)$, $CON(R_6, R_7)$, CH_2NO_2 , NO_2 , CH_2R_8 , CHR_9 , $-CH=N-OR_{10}$, $C=CH-R_5$, wherein R_5 is selected from the group consisting of H,

optionally substituted C_{1-12} alkyl, C_{3-12} cycloalkyl, aryl, heteroaryl; R_6 and R_7 are independently selected from the group consisting of H, optionally substituted C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy; R_8 and R_9 are independently selected from the group consisting of H, C_{1-6} alkyl, F, Cl, Br, C_{1-12} alkyl substituted with one or more of F, Cl, Br, I, OR_4 , SR_4 , wherein R_4 is the same as defined before, $N(R_6, R_7)$, R_{10} is selected from the group consisting of H, optionally substituted C_{1-12} alkyl, C_{3-12} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkyl, aryl, heteroaryl except $W = (CO)$, Q and P = H.


Ring C in Formula II is 6-8 membered or of larger size and the larger rings have either two or three carbons between each nitrogen atom, comprising of




and may be bridged to form a bicyclic system as shown below,




ring C is optionally substituted by Y and Z with alkyl groups, cycloalkyl groups, fluoro group, carboxylic and corresponding esters, amides, substituted alkyls or bridging alkyl groups are as shown below:



six membered ring C with $X = -CH-(NHR_{11})$, (wherein R_{11} is the same as defined earlier) is selected from the group consisting of the following rings;




wherein $M = \text{Sulphur}$ is shown by compounds of Formula III,




FORMULA III

wherein P, Q, U, V, X, Y, Z, W and n in Formula III are the same as previously defined, wherein the process comprising reacting a compound of Formula V



FORMULA V

with a compound of Formula VI



FORMULA VI

wherein P, Q, R₁₂, Y, Z, G, n, U and V are the same as defined earlier.


11. A process for preparing a compound of Formula II as claimed in claim 10, in a suitable solvent selected from the group consisting of dimethylformamide, dimethylacetamide, ethanol or ethylene glycol at a suitable temperature in the range of -70°C to 180°C in the presence of a suitable base selected from the group consisting of triethyl amine, diisopropyl amine, potassium carbonate and sodium bicarbonate.

12. A process of preparing a compound of Formula II as claimed in claim 10 wherein Formula VI is furalehyde and reductive alkylation of the amine of Formula V is performed with a reducing agent.

13. A process for preparing a compound of Formula II as claimed in claim 10 wherein Formula VI is furoic acid.

14. A process for preparing a compound of Formula II as claimed in claim 10 wherein the compounds of Formula II having carbonyl link are prepared by reacting heteroaromatic compound of the Formula VI including N- methyl pyrrole with the intermediate amine of Formula V in the presence of triphosgene or phosgene and carbonyl linkers are introduced between heteroaromatic compound comprising reacting 3- bromothiophene and amine of Formula V with carbon monoxide and the catalyst is selected from the group consisting of Pd (PPh₃)₂Cl₂ and extended chain pyrroles having dicarbonyl linkers are obtained by treatment of oxalyl chloride and amine of the Formula V.

15. A process for preparing a compound of Formula VIII



FORMULA VIII

wherein

n is an integer in the range from 0 to 3;

X is CH, CH-S, CH-O and N;

Y and Z are independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₃₋₁₂ cycloalkyl, C₀₋₃ bridging group;


5 **U and V** are independently selected from the group consisting of optionally substituted C₁₋₆ alkyl, F, Cl, Br, C₁₋₁₂ alkyl substituted with one or more of F, Cl, Br, I, preferably U and V are hydrogen or fluoro;

10 **W** is selected from the group consisting of CH₂, CO, CH₂NH, -NHCH₂, -CH₂NHCH₂, -CH₂-N(R₁₁)CH₂-, CH₂(R₁₁)N-, CH(R₁₁), S, CH₂(CO), NH wherein R₁₁ is optionally substituted with C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl, aryl, heteroaryl;

15 **Q and P** are independently selected from the group consisting of -CN, COR₅, COOR₅, N(R₆, R₇), CON(R₆, R₇), CH₂NO₂, NO₂, CH₂R₈, CHR₉, -CH=N-OR₁₀, C=CH-R₅, wherein R₅ is selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, aryl, heteroaryl; R₆ and R₇ are independently selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₆ alkoxy; R₈ and R₉ are independently selected from the group consisting of H, C₁₋₆ alkyl, F, Cl, Br, C₁₋₁₂ alkyl substituted with one or more of F, Cl, Br, I, OR₄, SR₄, wherein R₄ is the same as defined before, N(R₆, R₇), R₁₀ is selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl, aryl, heteroaryl except W= (CO), Q and P =H;


20

M = Sulphur is shown by compounds of Formula III



FORMULA III


and R_{15} is the same as Q defined earlier, comprising converting a compound of Formula VII



FORMULA VII


wherein in U, V, Y, Z, X, W, P, n and M are the same as defined earlier and R_{14} is any group which can be converted to group R_{15} in one to five steps.

16. A process for preparing a compound of Formula XI




FORMULA XI

($R_{16} = -CH_2F$ or $-CH_2F_2$) by reacting a compound of Formula IX



FORMULA IX


with sodium borohydride to produce a compound of Formula X



FORMULA X


and further reacting this compound with diethylamino sulfurtrifluoride to produce compound of Formula XI.

17. A process for preparing a compound of Formula XII



FORMULA XII


wherein $R_{17} = \text{---}N=OH$ which comprises reacting (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-formyl)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]-methyl] acetamide of Formula IX



FORMULA IX

with hydroxylamine.

18. A process for preparing a compound of Formula XII




5

FORMULA XII

wherein $R_{17} = \text{CH}_2=\text{N}-\text{NH}_2$ which comprises reacting (S)-N-[[3-[3-Fluoro-4[N-1-[4-{2-furyl-(5-hydrazone)-methyl}]]-piperazinyl]-phenyl]-2-oxo-5-oxazolidinyl]-methyl]acetamide with hydrazine hydrate.

19. A process for preparing a compound of Formula XII




10

FORMULA XII

wherein $R_{17} = \text{CH}_2=\text{N}-\text{O}-\text{C}(=\text{O})-\text{NH}-\text{C}_6\text{H}_4-\text{CH}_2\text{COOCH}_3$ which comprises reacting (S)-N-[[3-[3-Fluoro-4-[N-1-[4-(2-furyl-(5-aldoxime)methyl]] piperazinyl] phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide with isocyanate.

15

20. A process for preparing a compound of Formula XII




20


FORMULA XII

wherein $R_{17} = \text{CN}$ which comprises reacting (S)-N-[[3-[3-Fluoro-4-[N-1[4-{2-furyl(5-cyano)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide with triflic anhydride and triethylamine.


21. A process for preparing a compound of Formula XII




FORMULA XII

- 10 wherein $R_{17} =$  which comprises reacting (S)-N-[[3-Fluoro-4-[N-1[5-(1,3-dioxane)-2-furylmethyl]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide with 1,3-propane diol and BF_3 etherate.


22. A process for the preparation of the compound of Formula XIV



FORMULA XIV

wherein $R_{18} =$ 


- 20 which comprises reacting (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-formyl)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl] acetamide of Formula IX



FORMULA IX

5 with Ag_2O to produce (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-carboxy)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl] acetamide of Formula XIII followed by reacting (S)-N-[[3-Fluoro-4-[N-1[4-(2-furyl- (5-carboxyethyl)methyl)piperazinyl] phenyl]- 2-oxo-5-oxazolidinyl]methyl] acetamide of


Formula XIII




FORMULA XIII

with aqueous ammonia to produce Formula XIV.


23. A process for the preparation of the compound of Formula XIV



FORMULA XIV




which comprises reacting (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-formyl)-methyl}] piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl] acetamide of Formula IX



FORMULA IX


5 with Ag_2O to produce (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-carboxy)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl] acetamide of Formula XIII followed by reacting (S)-N-[[3-Fluoro-4-[N-1[4-(2-furyl- (5-carboxyethyl)methyl)piperazinyl] phenyl]- 2-oxo-5-oxazolidinyl]methyl] acetamide of Formula XIII



FORMULA XIII

with thionyl chloride to produce Formula XIV.


15 24. A process for the preparation of the compound of Formula XIV



FORMULA XIV


20 wherein $\text{R}_{18} =$

which comprises reacting (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-formyl)-methyl}] piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl] acetamide of Formula IX



FORMULA IX

5 with Ag_2O to produce (S)-N-[[3-Fluoro-4-[N-1[4-{2-furyl(5-carboxy)methyl}]piperazinyl]phenyl]-2-oxo-5-oxazolidinyl]methyl] acetamide of Formula XIII followed by reacting (S)-N-[[3-Fluoro-4-[N-1[4-(2-furyl- (5-carboxy-ethyl)methyl)piperazinyl] phenyl]- 2-oxo-5-oxazolidinyl]methyl] acetamide of Formula XIII



FORMULA XIII

with morpholine in the presence of oxalyl chloride to produce Formula XIV.